

Antiderivatives

William M. Faucette

University of West Georgia

Outline

- 1 General Instructions
- 2 Antiderivatives
- 3 Notation for Antiderivatives
- 4 Constant of Integration
- 5 Examples
- 6 Basic Antidifferentiation Formulas
- 7 Initial Value Problems and Differential Equations
- 8 Example
- 9 Indefinite Integrals
- 10 Examples

Antiderivatives

- As usual, you should read section 4.8 in the online textbook.
- This slideshow will give an overview and an explanation of the important concepts in the book.
- This slideshow will also include a limited number of examples.
- The main purpose of this slideshow is to give an extended explanation and clarification of the material in the text.

Antiderivatives

So far, we have been starting with a function $f(x)$, taking its derivative, and producing $f'(x)$.

Now, we reverse the process. We will be given $f'(x)$ and produce the function $f(x)$. The function $f(x)$ is the **antiderivative** of $f'(x)$.

Notation for Antiderivatives

If we are given a function $f(x)$, its antiderivative is denoted

$$F(x) = \int f(x) dx$$

or

$$F(x) = \int f.$$

The antiderivative is also called the **indefinite integral**.

Constant of Integration

Recall from the section the Mean Value Theorem and its corollary that if $f'(x) = g'(x)$, then $f(x) = g(x) + C$, for some constant C . This means that a given function doesn't have one antiderivative, but has a family of antiderivatives, all differing by a constant.

For example, the derivative of x^2 is $2x$. But the derivative of $x^2 + 3$, $x^2 - 6$, $x^2 + 47$, $x^2 + e - \pi$ is also $2x$.

We write this as

$$\int 2x \, dx = x^2 + C.$$

The C here is called the **constant of integration**.

Examples

- $\int 1 dx = x + C$ because $(x)' = 1$.
- $\int 2x dx = x^2 + C$ because $(x^2)' = 2x$.
- $\int \cos x dx = \sin x + C$ because $(\sin x)' = \cos x$.
- $\int 2^x \ln 2 dx = 2^x + C$ because $(2^x)' = 2^x \ln 2$.
- $\int \frac{1}{1+x^2} dx = \arctan x + C$ because $(\arctan x)' = \frac{1}{1+x^2}$.

The bad news is that this means you have to know the derivatives of functions to be able to find the antiderivatives.

Basic Antidifferentiation Formulas

$$\int \sin x \, dx = -\cos x + C$$

$$\int \cos x \, dx = \sin x + C$$

$$\int \sec^2 x \, dx = \tan x + C$$

$$\int \csc^2 x \, dx = -\cot x + C$$

$$\int \sec x \tan x \, dx = \sec x + C$$

$$\int \csc x \cot x \, dx = -\csc x + C$$

$$\int k \, dx = kx + C, \text{ for } k \text{ a constant}$$

$$\int x^n \, dx = \frac{x^{n+1}}{n+1} + C, \text{ for } n \neq -1$$

$$\int \frac{1}{x} \, dx = \ln|x| + C$$

Basic Antidifferentiation Formulas

$$\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + C$$

$$\int -\frac{1}{\sqrt{1-x^2}} dx = \arccos x + C$$

$$\int \frac{1}{1+x^2} dx = \arctan x + C$$

$$\int -\frac{1}{1+x^2} dx = \operatorname{arccot} x + C$$

$$\int \frac{1}{|x|\sqrt{x^2-1}} dx = \operatorname{arcsec} x + C$$

$$\int -\frac{1}{|x|\sqrt{x^2-1}} dx = \operatorname{arccsc} x + C.$$

$$\int a^x dx = \frac{a^x}{\ln a} + C, \text{ for } a > 0, a \neq 1$$

$$\int e^x dx = e^x + C,$$

Initial Value Problems and Differential Equations

An equation of the form $\frac{dy}{dx} = f(x)$ is called a **differential equation**. The goal here is to find the function y explicitly as a function of x .

If you're also given an initial condition $y(x_0) = y_0$, then the differential equation together with the initial condition is called an **initial value problem**.

You solve a differential equation of this form by taking the antiderivative. The initial condition then tells you the value of the constant of integration.

Example 1

Example

Solve the initial value problem

$$\frac{dy}{dx} = 9x^2 - 4x + 5, \quad y(-1) = 0.$$

Solution

$$\begin{aligned} y &= \int 9x^2 - 4x + 5 \, dx \\ &= 3x^3 - 2x^2 + 5x + C. \end{aligned}$$

Example 1

Solution

We substitute the initial value to find the value of C .

$$\begin{aligned}y &= 3x^3 - 2x^2 + 5x + C \\0 &= 3(-1)^3 - 2(-1)^2 + 5(-1) + C \\&= -3 - 2 - 5 + C \\C &= 10.\end{aligned}$$

So, the solution to the initial value problem is

$$y = 3x^3 - 2x^2 + 5x + 10.$$

Indefinite Integrals

Definition

The collection of all antiderivatives of f is called the indefinite integral of f with respect to x , and is denoted by

$$\int f(x) dx.$$

The symbol \int is an **integral sign**. The function f is the **integrand** of the integral, and x is the **variable of integration**.

Examples

Examples

$$\int (5 - 6x) dx = 5x - 3x^2 + C.$$

$$\int x^{-5/4} = \frac{x^{-1/4}}{-1/4} + C = -4x^{-1/4} + C.$$

$$\begin{aligned}\int (\sqrt{x} + \sqrt[3]{x}) dx &= \int (x^{1/2} + x^{1/3}) dx \\ &= \frac{x^{3/2}}{3/2} + \frac{x^{4/3}}{4/3} + C \\ &= \frac{2}{3}x^{3/2} + \frac{3}{4}x^{4/3} + C.\end{aligned}$$